The slide features a decorative layout with blue lines and corner markers. A vertical line on the left and a horizontal line at the top intersect at a small circle in the top-left corner. Another horizontal line is positioned below the title. A vertical line on the right and a horizontal line at the bottom intersect at a small circle in the bottom-right corner.

Heavy Ion Interactions

7th FLUKA Course
NEA Paris, Sept.29-Oct.3, 2008

Overview

The models

DPMJET
RQMD
BME

Input options

Beam definition
Transport thresholds

Heavy ion interaction models in FLUKA - 1

$E > 5 \text{ GeV/n}$

Dual Parton Model (DPM)

DPMJET-III (original code by R.Engel, J.Ranft and S.Roesler,
FLUKA-implemenation by T.Empl *et al.*)

$0.1 \text{ GeV/n} < E < 5 \text{ GeV/n}$

Relativistic Quantum Molecular Dynamics Model (RQMD)

RQMD-2.4 (original code by H.Sorge *et al.*,
FLUKA-implementation by A.Ferrari *et al.*)


$E < 0.1 \text{ GeV/n}$

Boltzmann Master Equation (BME) theory

BME (original code by E.Gadioli *et al.*,
FLUKA-implementation by F.Cerutti *et al.*)

Heavy ion interaction models in FLUKA - 2

Program flow:



DPMJET

$E > 5 \text{ GeV/n}$

Dual Parton Model (DPM)

DPMJET-III (original code by R.Engel, J.Ranft and S.Roesler,
FLUKA-implemenation by T.Empl *et al.*)

$0.1 \text{ GeV/n} < E < 5 \text{ GeV/n}$

Relativistic Quantum Molecular Dynamics Model (RQMD)

RQMD-2.4 (original code by H.Sorge *et al.*,
FLUKA-implementation by F.Cerutti *et al.*)

$E < 0.1 \text{ GeV/n}$

Boltzmann Master Equation (BME) theory

BME (original code by E.Gadioli *et al.*,
FLUKA-implementation by F.Cerutti *et al.*)

DPMJET - *Overview*

DPMJET = Dual Parton Model and JETs

Monte Carlo **event-generator** for the simulation of high-energy hadronic interactions

- hadron-hadron collisions,
hadron-nucleus collisions,
nucleus-nucleus collisions,
photoproduction off nuclei (only Version III)
- energy range: 5 GeV/nucleon – 10^{11} GeV/nucleon
- programming language: Fortran77
- size of the code (Version III): about 90.000 lines
- authors: J. Ranft (Version II),
R.Engel, J. Ranft, S. Roesler (Version III)

DPMJET - *Main steps of a high energy interaction*

1. Interaction of high-energy nuclei

- individual nucleon-nucleon scatterings
- formation of »strings« between valence and sea partons (quarks, gluons)



2. Hadronization process

- creation of hadrons / resonances




3. Intranuclear cascade

- low-energy interactions of hadrons in spectator nuclei



4. Fragmentation of excited spectator nuclei

- evaporation of light fragments (e.g., p, n, d, ^3H , ^3He , ^4He ,...),
- fragmentation, fission
- production of residual nuclei



DPMJET - *The Gribov-Glauber formalism*

Fundamental idea: nucleus-nucleus collision expressed in terms of individual nucleon-nucleon interactions

- nucleus-nucleus

- total cross section

$$\sigma_{AB}^{tot}(s) = 4 \int d^2 \vec{B} \mathfrak{I} [A_{AB}(s, \vec{B})]$$

- elastic cross section

$$\sigma_{AB}^{el}(s) = 4 \int d^2 \vec{B} \left| A_{AB}(s, \vec{B}) \right|^2$$

- scattering amplitude

$$A_{AB} = \frac{i}{2} \left[1 - \exp(\chi_{AB}) \right]$$

eikonal function

$$\chi_{AB} = \sum_{k,l} \chi_{N_k N_l}$$

- nucleon-nucleon

- scattering amplitude

$$a_{N_k N_l} = \frac{i}{2} \left[1 - \exp(\chi_{N_k N_l}) \right]$$

DPMJET - *Intranuclear cascade and fragmentation*

- nuclear model :
 - Fermi-gas of nucleons in potential well
 - nuclear densities: shell model ($A \leq 18$)
Wood-Saxon ($A > 18$)
- hadrons are followed in space and time on straight trajectories
- hadrons may re-interact after certain formation-time
(assume local nuclear density and vacuum cross sections)
- calculation of excitation energies (in AA interactions for both spectators)


Note: DPMJET has its own implementation of intranuclear cascades. It is similar in its fundamental ideas to that one of FLUKA but is much more simplified.

- fragmentation by
 - nuclear evaporation
 - Fermi-breakup
 - high-energy fission
 - γ -deexcitation
- } FLUKA

DPMJET - Comparison to data (hadron-hadron)




proton - proton

proton - proton, $E_{\text{lab}} = 200\text{GeV}$







	Exp.	DPMJET-III
charged	7.69 ± 0.06	7.64
neg.	2.85 ± 0.03	2.82
p	1.34 ± 0.15	1.26
n	0.61 ± 0.30	0.66
π^+	3.22 ± 0.12	3.20
π^-	2.62 ± 0.06	2.55
K^+	0.28 ± 0.06	0.30
K^-	0.18 ± 0.05	0.20
Λ	0.096 ± 0.01	0.10
$\bar{\Lambda}$	0.0136 ± 0.004	0.0105

DPMJET - Comparison to data (hadron-nucleus)




	Exp.	DPMJET-III
14.6 GeV p Al	1.57 ± 0.23	1.52
p Au	2.15 ± 0.33	1.92
200 GeV p S	5.0 ± 0.2	4.98
p Xe	6.84 ± 0.13	6.67
360 GeV p Al	6.8 ± 0.6	5.87
p Au	8.9 ± 0.4	8.77

DPMJET - Comparison to data (nucleus-nucleus)



DPMJET - Comparison to data (nucleus-nucleus)





Pseudorapidity distribution of charged hadrons produced in minimum bias d-Au and p-p collisions at a c.m. energy of 200GeV/A.

Exp. data: BRAHMS- and PHOBOS-Collaborations

J.Ranft, in Proceedings of the Hadronic Shower Simulation Workshop, CP896, Batavia, Illinois (USA), 6-8 September 2006

DPMJET - Comparison to data (nucleus-nucleus)




Pseudorapidity distribution of charged hadrons produced in Au-Au collisions at a c.m. energy of 130 GeV/A (left) and 200 GeV/A (right) for different ranges of centralities.

Exp. data: PHOBOS-Collaboration

J.Ranft, in Proceedings of the Hadronic Shower Simulation Workshop, CP896, Batavia, Illinois (USA), 6-8 September 2006


DPMJET - Interface to FLUKA



DPMJET - Interface to FLUKA


Examples for pre-initialised data:

Inelastic cross sections




$$E_{\text{Lab}} = 6.3 \times 10^9 \text{ GeV/nucleon}$$

Impact parameter distribution



e.g., for highest and lowest energy at one fixed projectile-target configuration

DPMJET - FLUKA benchmarks



Fragment charge cross sections for
158GeV/n Pb ions on various targets.
FLUKA: solid histogram(total)
dashed histogram (em diss.)
Exp. data: symbols
NPA662, 207 (2000),
NPA707, 513 (2002),
C.Scheidenberger *et al.* PRC

RQMD

$E > 5 \text{ GeV/n}$

Dual Parton Model (DPM)
DPMJET-III (original code by R.Engel, J.Ranft and S.Roesler,
FLUKA-implementation by T.Empl *et al.*)

$0.1 \text{ GeV/n} < E < 5 \text{ GeV/n}$

Relativistic Quantum Molecular Dynamics Model (RQMD)
RQMD-2.4 (original code by H.Sorge *et al.*,
FLUKA-implementation by A.Ferrari *et al.*)

$E < 0.1 \text{ GeV/n}$

Boltzmann Master Equation (BME) theory
BME (original code by E.Gadioli *et al.*,
FLUKA-implementation by F.Cerutti *et al.*)

RQMD - *References*

interface to a suitably modified **RQMD model**

RQMD-2.4 (H. Sorge, 1998) was successfully applied
to relativistic A-A particle production over a wide energy range

[H. Sorge, Phys. Rev. **C 52**, 3291 (1995);

H. Sorge, H. Stöcker, and W. Greiner, Ann. Phys. **192**, 266 (1989)

and Nucl. Phys. **A 498**, 567c (1989)]

RQMD - *The original code*

The RQMD-2.4 code

INITIAL CONDITION two Fermi gases (projectile and target)

$$\text{Fermi momentum } p_{F0} = \hbar \left(3\pi^2 \frac{A}{2V} \right)^{1/3} \quad V = (4/3) \pi (r_0 A^{1/3})^3 \quad r_0 = 1.12 \text{ fm} \Rightarrow \rho = 0.17 \frac{\text{nucl.}}{\text{fm}^3}$$

$$\text{nucleon momentum } \boxed{p = p_{F0} \left(\frac{\rho(r)}{\rho_0} \right)^{1/3} \epsilon^{1/3}} \quad \epsilon \in [0, 1] \text{ random}$$
$$\phi = 2\pi\epsilon \qquad \cos \theta = 1 - 2\epsilon$$

$$\begin{aligned} p_x &= p \sin \theta \cos \phi & - (\sum p_x) / A \\ p_y &= p \sin \theta \sin \phi & - (\sum p_y) / A \\ p_z &= p \cos \theta & - (\sum p_z) / A \end{aligned} \qquad \text{so } \sum p_x = \sum p_y = \sum p_z = 0$$

FINAL STATE


- (p^0, p_x, p_y, p_z) for nucleons (and produced particles) in the LAB frame
- the spectators are marked
- no residue and fragment identification
- energy non-conservation issues, particularly when run in full QMD mode

RQMD - The interfaced code

Implemented developments


- construct the **projectile- and target-like** nuclei by gathering *spectator* nucleons,

$$\text{assuming } E_{pL}^* = \sum_{pa} \rho E_h \quad (TL)$$



$$\rho(r) \propto \left(1 + \exp\left(\frac{r-R}{a}\right)\right)^{-1}$$

$$R = 1.19 A^{1/3} - 1.61 A^{-1/3} \text{ fm} \quad a = 0.52 \text{ fm}$$




$$E_h = \frac{1}{2m_n} \left\{ \left[p_{F0} (\rho(r)/\rho_0)^{1/3} \right]^2 - p^2 \right\}$$

$r, \rho(t = 0)$

- fix the remaining energy-momentum conservation issues taking into account **experimental binding energies**
- use the FLUKA evaporation/fission/fragmentation module


RQMD - FLUKA benchmarks

Double differential neutron yield



400 MeV/n



on *thick*
Al targets



exp. data from T. Kurosawa *et al.*, Phys. Rev. C **62**, 044615 (2000)

RQMD - FLUKA benchmarks


Fragment charge cross sections



exp. data from C. Zeitlin *et al.*, Phys. Rev. **C 56**, 388 (1997),
J.R. Cummings *et al.*, Phys. Rev. **C 42**, 2508 (1990) (at 1.5 GeV/n),
G.D. Westfall *et al.*, Phys. Rev. **C 19**, 1309 (1979) (at 1.88 GeV/n)

RQMD - FLUKA benchmarks

Isotopic distributions of fragmentation products



fission products excluded
like in the experimental
analysis

exp. data (stars) from J. Benlliure, P. Armbruster *et al.*, Eur. Phys. J A 2, 193 (1998)

7th FLUKA Course, NEA Paris, Sept.29–Oct.3, 2008

BME

$E > 5 \text{ GeV/n}$

Dual Parton Model (DPM)

DPMJET-III (original code by R.Engel, J.Ranft and S.Roesler,
FLUKA-implemenation by T.Empl *et al.*)

$0.1 \text{ GeV/n} < E < 5 \text{ GeV/n}$

Relativistic Quantum Molecular Dynamics Model (RQMD)

RQMD-2.4 (original code by H.Sorge *et al.*,
FLUKA-implementation by A.Ferrari *et al.*)

$E < 0.1 \text{ GeV/n}$

Boltzmann Master Equation (BME) theory

BME (original code by E.Gadioli *et al.*,
FLUKA-implementation by F.Cerutti *et al.*)

BME - *References*

interface to a Monte Carlo code
founded on the BME theory (E. Gadioli et al.)

[M. Cavinato *et al.*, Nucl. Phys. **A 679**, 753 (2001),

M. Cavinato *et al.*, Phys. Lett. **B 382**, 1 (1996)]

BME - *The interfaced code*

two different reaction paths have been adopted:

1. COMPLETE FUSION

$$P_{CF} = \sigma_{CF} / \sigma_R$$

pre-equilibrium

according to the BME theory

2. PERIPHERAL COLLISION

$$P = 1 - P_{CF}$$

three body mechanism

pickup/stripping (for asymmetric systems at low b)

inelastic scattering (at high b)

FLUKA evaporation


1. In order to get the multiplicities of the pre-equilibrium particles and their double differential spectra, the BME theory is applied to a few representative systems at different bombarding energies and the results are parameterized.
2. The complete fusion cross section decreases with increasing bombarding energy. We integrate the nuclear densities of the projectile and the target over their overlapping region, as a function of the impact parameter, and obtain a preferentially excited "middle source" and two fragments (projectile- and target-like). The kinematics is suggested by break-up studies.

BME - The database for the pre-equilibrium emissions

$^{16}\text{O} + ^6\text{Li}, ^8\text{Li}, ^8\text{B}, ^{10}\text{B}, ^{12}\text{C}, ^{14}\text{N}, ^{16}\text{O}, ^{19}\text{F}, ^{20}\text{Ne}, ^{24}\text{Mg}, ^{27}\text{Al}$


$^{12}\text{C} + ^8\text{Li}, ^8\text{B}, ^{12}\text{C}, ^{27}\text{Al}$

@ 12, 30, 50, 70, 100 MeV/n



total multiplicity

$$M = P_1 E_{nucl} - P_2 E_{nucl}^2$$



energy spectra

$$\frac{d^2 M}{(dE d\Omega)} = E^{P_0(\theta)} \exp(-P_1(\theta) - P_2(\theta)E)$$

BME - Theoretical framework


Calculation of preequilibrium for the composite nucleus

proton and neutron momentum spaces divided into **bins**

$$\left\{ (p_x, p_y, p_z) : p_z \in [p_{zi}, p_{zi} + \Delta p_z), \varepsilon = (2m)^{-1} (p_x^2 + p_y^2 + p_z^2) \in [\varepsilon_i, \varepsilon_i + \Delta\varepsilon) \right\}$$

(Z is the beam direction)

of volume $2\pi m \Delta\varepsilon \Delta p_z$



BME - Theoretical framework

The BME system

$$N_i = n_i g_i$$

nucleon number
number of states in bin i

|
|

occupation probability

$$\begin{aligned}
 \frac{d(n_i^\pi g_i^\pi)}{dt} = & \sum_{jlm} [\omega_{lm \rightarrow ij}^{\pi\pi} g_i^\pi n_l^\pi g_m^\pi n_m^\pi (1 - n_i^\pi)(1 - n_j^\pi) \\
 & - \omega_{ij \rightarrow lm}^{\pi\pi} g_i^\pi n_i^\pi g_j^\pi n_j^\pi (1 - n_l^\pi)(1 - n_m^\pi)] \\
 + & \sum_{jlm} [\omega_{lm \rightarrow ij}^{\pi\nu} g_i^\pi n_l^\pi g_m^\nu n_m^\nu (1 - n_i^\pi)(1 - n_j^\nu) \\
 & - \omega_{ij \rightarrow lm}^{\pi\nu} g_i^\pi n_i^\pi g_j^\nu n_j^\nu (1 - n_l^\pi)(1 - n_m^\nu)] \\
 - & n_i^\pi g_i^\pi \omega_{i \rightarrow i'}^\pi g_{i'}^\pi \delta(\epsilon_i^\pi - \epsilon_{i'}^\pi - \epsilon_F^\pi - B^\pi) - \frac{dD_i^\pi}{dt}
 \end{aligned}$$

BME - Theoretical framework

Multiplicity spectra

of emitted nucleons


$$\frac{d^2 M(\varepsilon', \theta)}{d\varepsilon' d\Omega} = \frac{1}{2\pi \sin \theta} \int_0^{t_{eq}} n(\varepsilon, \theta, t) \frac{\sigma_{inv} V}{V} \rho(\varepsilon', \theta) dt$$

of a cluster c

$$\frac{d^2 M_c(E'_c, \theta_c)}{dE'_c d\Omega} = \frac{R_c}{2\pi \sin \theta} \int_0^{t_{eq}} N_c(E_c, \theta_c, t) \frac{\sigma_{inv,c} V_c}{V} \rho_c(E'_c, \theta_c) dt$$

$$N_c(E_c, \theta_c, t) = \prod_i (n_i^\pi(\varepsilon, \theta, t))^{P_i(E_c, \theta_c) Z_c} \cdot \prod_i (n_i^\nu(\varepsilon, \theta, t))^{P_i(E_c, \theta_c) N_c}$$

joint probability



BME - Peripheral collisions

i. selection of the *impact parameter* b

ii. kinematics determination

θ_{PL} , θ_{TL} chosen according to $[d\sigma/d\Omega]_{cm} \sim \exp(-k\theta_{cm})$



θ_{MS} momentum conservation

p_{PL} , p_{TL} chosen according to a given energy loss distribution

p_{MS} momentum conservation

ϕ_{PL} free

ϕ_{TL} , ϕ_{MS} same reaction plane



iii. excitation energy sharing

$$E_{MS}^* = (A_{MS}/A_{tot}) E_{tot}^* \sum_{n=0}^k (1 - A_{MS}/A_{tot})^n$$


$$E_{PL}^* = f(A_{PL}, A_{TL}) (E_{tot}^* - E_{MS}^*)$$

$$E_{TL}^* = (E_{tot}^* - E_{MS}^* - E_{PL}^*)$$

BME - Benchmarking

Double differential neutron yields from thick targets

exp. data from T. Kurosawa, N. Nakao, T. Nakamura *et al.*, Nucl. Sci. Eng. 132, 30-57 (1999)







100 MeV/n ^{12}C ions on C target

BME - Benchmarking

Fragment production

exp. data from H. Ryde, Physica Scripta T5, 114-117 (1983)



86 MeV/n ^{12}C ions on ^{12}C

$2^\circ < \theta_{\text{LAB}} < 22^\circ$ $T_{\text{LAB}} > 10\text{MeV/n}$

Input options - 1

a) define momentum / energy

```
BEAM          -10.0      0.0      0.0      0.0      0.0      0.0HEAVYION
```

WHAT(1) > 0.0 : average beam momentum (GeV/c)
< 0.0 : average beam kinetic energy (GeV)

WHAT(2) beam momentum spread (GeV/c)

Note: for SDUM = HEAVYION units per nucleon
for SDUM = 4-HELIUM, etc. units per nucleus

WHAT(3)-WHAT(6) (as for any other particle)

SDUM = HEAVYION

also 4-HELIUM alpha
3-HELIUM 3-helium
TRITON tritium
DEUTERON deuterium

Input options - 2

b) define charge and mass (*required for* BEAM/SDUM=HEAVYION)

```
HI-PROPE          79.0      197.0      0.0      0.0      0.0      0.0
```

WHAT(1) = Atomic number Z of the heavy ion, Default: 6.0

WHAT(2) = Mass number A of the heavy ion, Default: 12.0

WHAT(3) = if < 0 isomeric state of the heavy ion

c) switch on heavy ion interactions

```
EVENTYPE          0.0      0.0      2.0      0.0      0.0      0.0DPMJET
```

Note: Don't forget to link the DPMJET/RQMD modules.
either using FLAIR or

```
$FLUPRO/flutil/ldpmqmd -m fluka -o <executable name>
```

Input options - 3

Notes:

- The transport momentum threshold for ions ($p_{th,HI}$) is fixed to that of alphas ($p_{th,\alpha}$)

$$p_{th,HI} = p_{th,\alpha} \times m_{HI}/m_{\alpha} \quad (GeV/c)$$

- If the transport momentum threshold for alphas is not explicitly defined with a PART-THR card (*requiring GeV and not GeV per nucleon*) it is fixed to that of protons ($p_{th,p}$).

$$p_{th,\alpha} = p_{th,p} \times m_{\alpha}/m_p \quad (GeV/c)$$

- Unless the transport threshold for protons is defined with a PART-THR card it is set equal to 10 MeV if `DEFAULTS=NEW-DEFA`.
- When the energy of an ion becomes lower than the transport threshold, and if such threshold is lower than 100 MeV/n, the ion is not stopped, but it is ranged out to rest

Input options - 4

Electromagnetic dissociation

PHYSICS	0.0	0.0	0.0	0.0	0.0	0.0EM-DISSO
---------	-----	-----	-----	-----	-----	-------------

WHAT(1) : flag for activating ion electromagnetic-dissociation

- =< -1.0 : resets to default (no em-dissociation)
- = 0.0 : ignored
- = 1.0 : (default) no em-dissociation
- = 2.0 : projectile and target em-dissociation activated
- = 3.0 : projectile only em-dissociation activated
- = 4.0 : target only em-dissociation activated

WHAT(2)-WHAT(6): not used