

The FLUKA Code


An Introduction to FLUKA: a multipurpose Interaction and Transport MC code

Beginners FLUKA Course

FLUKA

Main authors: A. Fassò, A. Ferrari, J. Ranft, P.R. Sala

Contributing authors: G.Battistoni, F.Cerutti, T.Empl, M.V.Garzelli, M.Lantz, A. Mairani, V.Patera, S.Roesler, G. Smirnov, F.Sommerer, V.Vlachoudis



Developed and maintained under an INFN-CERN agreement Copyright 1989-2009 CERN and INFN

>2000 users

http://www.fluka.org

The FLUKA international Collaboration

M.Brugger, F. Cerutti, A. Ferrari, G. Lukasik, S. Roesler, G. Smirnov, F. Sommerer, C. Theis, S. Trovati, H. Vinke, V.Vlachoudis CERN



- A. Fassò, J. Vollaire SLAC, USA
- J. Ranft Univ. of Siegen, Germany
- G. Battistoni, F. Broggi, M. Campanella, P. Colleoni, E. Gadioli, A. Mairani, S. Muraro, P.R. Sala INFN & Univ. Milano, Italy
- L. Sarchiapone INFN Legnaro, Italy
- M. Carboni, C. D'Ambrosio, A. Ferrari, A. Mostacci, V. Patera, M. Pelliccioni, R. Villari INFN Frascati, Italy



- M.C. Morone Univ. Roma II, Italy
- A. Margiotta, M. Sioli INFN & Univ. Bologna, Italy
- K. Parodi DKFZ & HIT, Heidelberg, Germany
- A. Empl, L. Pinsky Univ. of Houston, USA
- K.T. Lee, T. Wilson, N. Zapp NASA-Houston, USA
- S. Rollet ARC Seibersdorf Research, Austria
- M. Lantz Riken, Japan



FLUKA Applications

Momentun The LHC Point 6 **Loss Regions** egions of high losses (e.g., Collimators,...) Point 7 Regions with low losses (e.g., due to residual gas) Point 2

0.8

9.0 it

Point 1


CMS

LHCb

LHC Dump

Betatron Cleaning

- Cosmic ray physics
- Neutrino physics
- Accelerator design (→ n_ToF, CNGS, LHC systems)
- Particle physics: calorimetry, tracking and detector simulation etc.
 - $(\rightarrow ALICE, ICARUS, ...)$
- ADS systems, waste transmutation, (\rightarrow "Energy amplifier", FEAT, TARC,...)
- Shielding design
- Dosimetry and radioprotection
- Space radiation
- Hadrontherapy
- **Neutronics**



The History

The early days

The beginning:

1962: Johannes Ranft (Leipzig) and Hans Geibel (CERN): Monte Carlo for high-energy proton beams

The name:

1970: study of event-by-event fluctuations in a NaI

calorimeter (FLUktuierende KAskade)

Early 70's to ≈1987: J. Ranft and coworkers (Leipzig University) with contributions from Helsinki University of Technology (J. Routti, P. Aarnio) and CERN (G.R. Stevenson, A. Fassò)

Link with EGS4 in 1986, later abandoned

The modern code: some dates

Since 1989: mostly INFN Milan (A. Ferrari, P.R. Sala): little or no remnants of older versions. Link with the past: J. Ranft and A. Fassò

1990: LAHET / MCNPX: high-energy hadronic FLUKA generator *No further update*

1993: G-FLUKA (the FLUKA hadronic package in GEANT3). *No further update*

1998: FLUGG, interface to GEANT4 geometry

2000: grant from NASA to develop heavy ion interactions and transport

2001: the INFN FLUKA Project

2003: official CERN-INFN collaboration to develop, maintain and distribute FLUKA

The FLUKA Code design - 1

- Sound and updated physics models
 - Based, as far as possible, on original and well-tested microscopic models
 - Optimized by comparing with experimental data at single interaction level: <u>"theory driven, benchmarked with data"</u>
 - Final predictions obtained with minimal free parameters fixed for all energies, targets and projectiles
 - Basic conservation laws fulfilled "a priori"
 - → Results in complex cases, as well as properties and scaling laws, arise naturally from the underlying physical models
 - → Predictivity where no experimental data are directly available

It is a "condensed history" MC code, with the possibility use of single instead of multiple scattering

The FLUKA Code design - 2

- Self-consistency
 - Full cross-talk between all components: hadronic, electromagnetic, neutrons, muons, heavy ions
 - Effort to achieve the same level of accuracy:
 - for each component
 - for all energies
 - Correlations preserved fully within interactions and among shower components
 - → FLUKA is NOT a toolkit! Its physical models are fully integrated

The Physics Content of FLUKA

- Nucleus-nucleus interactions 100 MeV/n 10000 TeV/n
 New model (BME, under development): from Coulomb Barrier
- Electromagnetic and µ interactions 1 keV 10000 TeV
- Hadron-hadron and hadron-nucleus interactions 0–10000 TeV
- Neutrino interactions (new DIS and RES generator!)
- Charged particle transport including all relevant processes
- Transport in magnetic field
- Neutron multigroup transport and interactions 0 20 MeV
 new library with 260 groups
- Analog calculations, or with variance reduction

Code complexity

- Inelastic h-N: ~ 72000 lines
- Cross sections (h-N and h-A), and elastic (h-N and h-A): ~32000 lines
- ➤ (G)INC and preequilibrium (PEANUT): ~114000 lines
- Evap./Fragm./Fission/Deexc.: ~27000 lines
- > v-N interactions: ~35000 lines
- A-A interactions:
 - ✓ FLUKA native (including BME): ~8000 lines
 - ✓ DPMJET-3: ~130000 lines
 - √ (modified) rQMD-2.4: ~42000 lines
- □ FLUKA in total (including transport, EM, geometry, scoring): ~680000 lines
- □ ... + ~20000 lines of ancillary off-line codes used for data pregeneration
- ☐ ... and ~30000 lines of post-processing codes

- General use: installation and run by a single user on a single machine
- Run can be managed also in computer farms (clusters)
- experimented under GRID

The FLUKA course: an Introduction

How:

This course is intended to provide users with the basic (and possibly more than basic!) knowledge of:

- a) The most relevant FLUKA instructions and options
- b) The physics models adopted in FLUKA
- c) The different scoring options embedded in FLUKA
- d) The different running options
- e) How to insert user code in FLUKA
- f) The tools to plot results
- g) The right approach to the existing documentation
- The procedures to overcome difficulties and problems and related debugging tools
- i) etc. etc.

Method:

There will be formal lectures but they will be followed much as possible by practical (simple) examples.

Emphasis will be given to practice.

If possible we shall try to transform your questions into cases of general interest.

A possible problem:

People here are not at the same level of FLUKA knowledge. There are those who already have some experience, maybe not negligible.

However we need to start from scratch.

We apologize to the experienced people and beg them to be patient: it's not excluded a-priori that they can learn something new also concerning the very basic elements!

A glimpse of FLUKA

The FLUKA version

FLUKA2008.n(x)(.m)

Major version

Minor version Patch
level

Since 2006 each version is going to be maintained for 2 years max.

In this course we are using FLUKA2008.3 (just released a few days ago)

The FLUKA license (it is not GPL):

- Standard download: binary library + user routines.
 - FLUKA can be used freely for scientific and academic purposes, ad-hoc agreement for commercial purposes
 - It cannot be used for weapon related applications
 - It is not permitted to redistribute the code (single user, single site)
 - User can add their own scoring, sources etc through a wide set of user routines, provided they don't modify the physics
 - Relevant references for each FLUKA version can be found in the documentation
- It is possible, by explicit signature of license, to download (!!! now from NEA as well !!!) the source for researchers of scientific/academic Institutions.
 - FLUKA cannot be copied, even in part, into other codes, or translated into another language without permission.
 - The user cannot publish results with modified code, unless explicit authorization is granted in advance.

Using FLUKA

Platform: Linux with g77, Linux with g95, Mac OSX with g9\$

Under test: Linux and Mac OSX (gfortran), Windows-Cygwin (g95)

The code can compiled/run only on with operating systems, compilers (and associated) options tested and approved by the development team


Standard Input:

- Command/options driven by "data cards" (ascii file) . Graphical interface is available!!!!
- Standard Geometry ("Combinatorial geometry"): input by "data cards"

Standard Output and Scoring:

- Apparently limited but highly flexible and powerful
- Output processing and plotting interface available

Examples of user interfaces



FLUKA Release Notes

Thi FLUKA release used in this course is the current official release version Fluka2008.3b.0. As it is obvious from the name, the major revision number of this release is 2008 and the minor revision number is 3. Patch b. Respin number is 0.

This release contains a few small fixes and a significant functionality improvement with respect to Fluka2008.3.7 (the latest respin of Fluka2008.3).

New functionality:

Compton scattering with full account for binding and orbital electron motion. Up to now FLUKA included two possibilities for the treatment of Compton scattering:

- 1) "naive" scattering on free electrons
- 2) Compton scattering corrected by an inelastic form factor, S(q,Z)

Now a third possibility has been added, where both binding effects and orbital motion of all electronic shells of all elements are accounted for. This is particularly relevant for low energy photons and/or heavy elements

FLUKA is also distributed by NEA

Starting from this release, the code will be distributed in parallel to the FLUKA web site by the Nuclear Energy Agency (NEA-OECD) Data Bank.

The NEA-OECD distribution (source included) will operate under the same license and conditions, and following the procedures specific of NEA-OECD.

FLUKA Release New Features


- New neutron cross section library below 20 MeV, including 260 neutron and 42 gamma groups: 31 neutron groups are thermal (1 in the previous library) (thanks to D. Ene, A. Fassò, A.Ferrari, G. Panini, F. Sommerer) . .
 - Please note that the new 260 group library is now the default one (even though the "old" 72 group one is still distributed).
- Heavy ion pair production
- New implementation of the BME model with vastly improved performances for peripheral collisions (BME is available on request)
- A new neutrino-nucleus event generator, including quasi-elastic, resonance, and deep-inelastic interactions (thanks to M. Lantz, P.R. Sala, G. Smirnov, G. Battistoni, and A. Ferrari)
- An enhanced version of the PEANUT event generator which should significantly improve residual nuclei predictions in the intermediate energy range, and more in general should further improve predicted particle spectra

FLUKA Release New Features

- New radioactive decay database, now including also conversion electron and Auger lines
- New generalized particles:
 - Dose (GeV/g) (DOSE, generalized particle id 228)
 - Dose equivalent (pSv) (DOSE-EQ, generalized particle id 240)
 - 1 MeV neutron Si equivalent fluence (for Silicon damage)
 (SI1MEVNE, generalized particle id 236)
 - High energy hadron fluence scoring (hadrons with energy larger than 20 MeV) (HADGT20M, generalized particle id 237)
- New option AUXSCORE
 - easy filtering of estimators (see scoring lecture)
 - choice of conversion coefficients for DOSE-EQ


Examples of FLUKA Applications

The TARC experiment at CERN:




The TARC experiment: neutron spectra


FLUKA + EA-MC (C.Rubbia et al.)




Applications – LHC collimation region





Applications – *LHC collimation region*



Applications – CNGS







Neutrino interactions


NunDIS is a new neutrino-nucleon Deep-Inelastic-Scattering event generator developed for FLUKA (M. Lantz, A. Ferrari, G. Battistoni, P. Sala, G. Smirnov)

- Designed to be fully integrated in FLUKA
 - → use the same hadronization routines as hadron-nucleon interactions
 - → all nuclear effects taken into account by PEANUT
- Energy range: from threshold to 10 TeV


NunRES is a generator of neutrino-nucleon resonant interactions

 No non-resonant background term, assuming that the non-resonant contribution comes from NunDIS

QUASI-ELASTIC interactions already implemented since 1997



(3D) Calculation of Atmospheric ν Flux



The first 3-D calculation of atmospheric neutrinos was done with FLUKA.


The enhancement in the horizontal direction, which cannot be predicted by a 1-D calculation, was fully unexpected, but is now generally acknowledged.


In the figure: angular distribution of v_{μ} , v_{μ} , v_{e} , v_{e} .

In red: 1-D calculation


Atmospheric muons simulations and comparison with exp. data

BESS 2001 balloon flight @ Ft. Summer






Application to ICARUS exp. at LNGS



Neutrons on the ER-2 plane at 21 km altitude




FLUKA calculations:

Roesler et al., Rad. Prot. Dosim. 98, 367 (2002)


Measurements: Goldhagen et al., NIM A476, 42 (2002)

Note one order of magnitude difference depending on latitude




Dosimetry Applications

Roesler et al., Rad. Prot. Dosim. 98, 367 (2002)



Ambient dose equivalent from neutrons at solar maximum on commercial flights from Seattle to Hamburg and from Frankfurt to Johannesburg.


Solid lines: FLUKA simulation



Experimental validation against measured Bragg curve in Proton and Carbon ion therapy

Protons (183 MeV/u) in Water

¹²C ions (400 MeV/u) in Water




Exp. Data (points) taken at HIT:

D. Schardt, P. Steidl, K. Parodi,

S. Brons et al.

Simulation: K. Parodi




Exp. Data (points) from Haettner et al, Rad. Prot. Dos. 2006 Simulation: A. Mairani, PhD Thesis, Pavia, 2007


Using the information from the patient CT in the MC I The Voxel Geometry

- FLUKA can embed voxel structures within its standard combinatorial geometry
- Transport through the voxels is optimized and efficient
- Raw CT-scan outputs can be imported

The GOLEM phantom enss et al,



Proton therapy: MC vs Focus/XiO for a Clivus Chordoma Patient at MGH




Parodi et al, JPCS 74, 2007

Prescribed dose: 1 GyE

 $MC: \sim 5.5 \cdot 10^6$ protons in 10 independent runs (11h each on Linux Cluster mostly using 2.2GHz Athlon processors)


Space radiation: GCR/SEP doses in

open space



- Capsule like enclosure with variable Al thickness
- Impinging uniform and isotropic GCR's and SEP "open space" spectra
- Scoring (each organ):
 - ✓ Organ Dose (GCR: Gy/day, SEP: Gy/event)
 - ✓ Organ Dose Equivalent (GCR: Sv/day, SEP: Sv/event), using the ICRP60 Q(LET) relationship
 - ✓ Effective Dose (whole body)
 using the newest ICRP
 recommendations (in press) (GCR:
 Sv/day, SEP: Sv/event)

GCR: open space doses after 1 g/cm² Al: Whole Body Dose Equivalent vs Effective Dose



The effective dose definition is clearly nonsense for relativistic heavy ions; in particular the \mathbf{w}_R =20 for \mathbf{a} 's and all heavy ions is largely overestimating their contribution

The FLUKA mailing lists

fluka-users@fluka.org

Users are automatically subscribed here when registering on the web site. It is used to communicate the availability of new versions, patches, etc.

fluka-discuss@fluka.org

Users are encouraged to subscribe at registration timed, but can uncheck the relevant box. It is used to have user-user and user-expert communication about problems, bugs, general inquiries about the code and its physics content

users are strongly encouraged to keep this subscription

